

Leggi attentamente il seguente brano e rispondi alle domande di comprensione.

online.zanichelli.it/chimicanatura

VERSO IL CLIL

How sweet it is!



SCHEDA VERSO IL CLIL



How sweet it is!


Did you think artificial sweeteners were a product of the post-World War II chemical industry? Not so –many of them have been around a long time, and several of the important ones were discovered quite by accident. In 1878, Ira Remsen was working late in his laboratory and realized he was about to miss a dinner with friends. In his haste to leave the lab, he forgot to wash his hands. Later at dinner he broke a piece of bread and tasted it only to discover that it was very sweet. The sweet taste had to be the chemical he had been working with in the lab. Back at the lab, he isolated saccharin- the first of the artificial sweeteners.

In 1937, Michael Sveda was smoking a cigarette in his laboratory (a very dangerous practice to say the least!). He touched the cigarette to his lips and was surprised by the exceedingly sweet taste. The chemical on his hands turned out to be cyclamate, which soon became a staple of the artificial sweetener industry.

In 1965, James Schlatter was researching antiulcer drugs for the

pharmaceutical firm G. D. Searle. In the course of his work, he accidentally ingested a small amount of a preparation and found to his surprise that it had an extremely sweet taste. He had discovered aspartame, a molecule consisting of two amino acids joined together. Since only very small quantities of aspartame are necessary to produce sweetness, it proved to be an excellent low-calorie artificial sweetener. More than 50 different molecules have a sweet taste, and it is difficult to find a single binding site that could interact with all of them.

Our taste receptors are composed of proteins that can form hydrogen bonds with other molecules. The proteins contain $-NH$ and $-OH$ groups (with hydrogen available to bond) as well as $C=O$ groups (providing oxygen for hydrogen bonding). «Sweet molecules» also contain H-bonding groups including $-OH$, $-NH_2$, and O or N . These molecules not only must have the proper atoms to form hydrogen bonds, they must also contain a hydrophobic region (repels H_2O). A new model for binding to a sweet-



ness receptor has been developed at Senomyx in La Jolla, California. The model shows four binding sites that can act independently. Small molecules bind to a pocket on a subunit as shown in the model. Large molecules (such as proteins) bind to a different site above one of the pockets.

- A Which was the first of artificial sweeteners?
- B How do taste receptors work?